



Research of hemoparasites in ticks collected from dogs resident in the municipality of Itu, São Paulo, SP

Tânia Regina Vieira de Carvalho¹; Paula Rocha Andrade²; Jonas Moraes-Filho^{1,2*}

¹Programa de Doutorado com ênfase em Saúde Única, Universidade Santo Amaro, São Paulo/SP, Brasil.

²Faculdade de Medicina Veterinária, Universidade Santo Amaro, São Paulo/SP, Brasil.

ABSTRACT

OBJECTIVE

The aim of this study was to report the occurrence of *Ehrlichia canis*, *Babesia canis vogeli* and *Rangelia vitalli* in ticks collected from dogs living in the city of Itu, São Paulo/SP, Brazil.

METHODS

DNA was extracted from 200 tick samples using the extraction kit PureLink Genomic DNA Kit (Invitrogen®) and real-time PCR was performed for the detection of *Ehrlichia canis*, *Rangelia vitalli* and *Babesia canis vogeli*.

RESULTS

We tested 200 ticks, 1/200 (0.5%) of the genus *Amblyomma aureolatum* and 199/200 (99.5%) of *Rhipicephalus sanguineus*. The results show an occurrence rate of positivity only in *R. sanguineus*, being 0.5% (1/200) for *E. canis*; 41% (82/100) for *B. c. vogeli*. No ticks were positive for *R. vitalli*.

CONCLUSIONS

The detection of *B. canis vogeli* and *E. canis* in ticks collected from dogs living in the municipality of Itu, São Paulo State, Brazil; show the dispersion of these pathogens in the country and the role of *R. sanguineus* s.l. as a vector of these pathogens cannot be neglected.

DESCRIPTORS

Ehrlichia canis, *Babesia canis vogeli*, *Rangelia vitalli*, Ticks.

Corresponding author:

Jonas Moraes-Filho.

Docente no Programa de Mestrado e Doutorado em Medicina e Bem-estar Animal e Saúde Única, Universidade Santo Amaro. R. Prof. Enéas de Siqueira Neto, 340 - Jardim das Imbuías, São Paulo - SP, Brasil

E-mail: jmfilho@prof.unisa.br

ORCID ID: <https://orcid.org/0000-0002-4734-9512>

Copyright: This is an open-access article distributed under the terms of the Creative Commons

Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided that the original author and source are credited.

INTRODUCTION

Of the approximately 1825 tick species described in the world, only about 10% are of direct public health importance, due to the possibility of parasitizing humans^{1,2}. Several other tick species that have never been described parasitizing humans assume indirect importance in public health, as they contribute to the enzootic maintenance of infectious agents in nature³.

Among the species with great importance for public and animal health, we can mention *Amblyomma sculptum* and *Amblyomma aureolatum*, vectors of the bacterium *Rickettsia rickettsii*, the causative agent of Brazilian Spotted Fever⁴; the last one also transmits the protozoan *Rangelia vitalli*, the pathogenic agent that causes the disease Canine Rangeliosis⁵. The tick *Rhipicephalus sanguineus* sensu lato (s.l.) is cosmopolitan and capable of transmitting microorganisms such as *Ehrlichia canis* and *Babesia canis vogeli* in domestic dogs⁶.

Canine Monocytic Ehrlichiosis (CME) is caused by *Ehrlichia canis*, an obligate intracellular bacterium found inside monocytes and macrophages of domestic dogs. It is a multisystem disease, with acute, subclinical or chronic clinical presentations. Transmission occurs in the blood meal of the infected *R. sanguineus* tick⁷.

Canine Babesiosis is an emerging disease of great importance in veterinary medicine due to its distribution, infectivity and pathogenesis⁸, whose etiological agent is intraerythrocyte protozoa of the genus *Babesia*⁹, species *Babesia canis*, in which it was subdivided into three subspecies: *Babesia canis canis*, *Babesia canis rossii* and *Babesia canis vogeli*⁹. In Brazil, the prevalent subspecies in dogs is *B. canis vogeli* and is distributed in tropical, subtropical and Mediterranean regions^{9,10}, present in both urban and rural environments¹¹, but mainly in urban and peripheral areas¹².

The protozoan *Rangelia vitalli* belongs to the Order *Piroplasmorida* and infects erythrocytes and endothelial cells of canids¹³. In a recent work carried out by Soares and collaborators⁵; it was reported that the species *A. aureolatum* is the only one that demonstrated vector competence for *R. vitalli*, as it was able to acquire and transmit the agent between domestic dogs.

Considering the risk for the occurrence of pathogens transmitted by these Ixodidae¹ in domestic dogs, it is essential to carry out research regarding the occurrence of *E. canis*, *B. canis vogeli* and *R. vitalli* in ticks in Brazil.

METHODS

Ticks were collected from 289 dogs that were screened at the Centro de Controle de Zoonoses de Itu, São Paulo, for pre-surgical evaluation of sterilization from March to November 2016. The ticks were preserved in flasks containing 70° alcohol. and duly identified according to Barros-Battesti¹⁴.

The study was approved by the Ethics Committee for the Use of Animals at Universidade Santo Amaro (CEUA 15/2018).

Ticks were processed individually. DNA extraction was performed with the Purelink Genomic DNA Extraction Kit (Invitrogen®), according to the manufacturer's instructions. The eluates obtained from DNA were properly identified and stored at -20° C for further molecular analysis.

Real-time PCR for: a) *Babesia canis vogeli* was performed using sense hsp70-F and antisense hsp70-R primers associated with a specific internal fluorogenic probe (5'-Hex/AGCGCCAG-GCCACCAAGGACGCT-3'-IABlkFQ), obtaining the amplification of a fragment of the hsp70 gene¹⁵; b) for *E. canis*, real-time PCR was performed using primers Dsb-321 and Dsb-671, in addition to the specific probe TaqMan (5'-AGCTAGTGCTGCTTG-GGCAACTTTGAGTGAA - 3') 5' FAM/BHQ - 1 3', obtaining an amplified nucleotide sequence of the dsb gene¹⁶ is shown; c)

for *R. vitalli* he used the oligonucleotide primers called sense Rv751-770 and antisense Rv930-91, in addition to a TaqMan probe [5'-6-FAM (CCT TAT CAA ATC ATT CTT C) MGB NFQ -3']. This pair of primers corresponds to the amplification of a fragment of the hsp70 gene¹³.

The reactions were performed in 96-well plates subjected to thermal variations corresponding to an initial cycle of 95° C for 5 minutes, followed by 40 cycles of 95° C for 15 seconds and 60° C for one-minute²⁸. Amplification, acquisition, and data analysis were performed using the multicolor detection system for Real-Time PCR (7500 Real-Time PCR Systems - Applied Biosystems, Foster City, CA, USA).

RESULTS

A total of 200 ticks were tested, 1/200 (0.5%) of the species of the genus *Amblyomma aureolatum* and 199/200 (99.5%) of *Rhipicephalus sanguineus*. The results show an occurrence rate of positivity only in *R. sanguineus*, being 0.5% (1/200) for *E. canis*; 41% (82/100) for *B. canis vogeli*. No ticks were positive for *R. vitalli*.

DISCUSSION

The largest number of ticks collected in the present work was of the species *R. sanguineus* (97%), a fact that was already expected, because they were collected from dogs that are their main hosts (64/65) and such arthropods have a nidicola habit with great proximity to dogs and humans¹².

Ticks of the species *A. aureolatum* can be found parasitizing dogs, with wild carnivores as their main natural hosts in a native area of the Atlantic Forest, the lowest percentage of the finding of such tick species in dogs in the present study.

There is a complex interaction between hosts, vectors, arthropods and pathogens and the low prevalence of *E. canis* in the present study may be related to the decrease in infected animals for the maintenance of the pathogen in a tick population, as there is no vertical transmission in the tick population *R. sanguineus*¹⁹.

Since *B. canis vogeli* causes a chronic and mild disease, it provides infected dogs for a longer time and consequently "food" for the ticks to maintain their infected populations for a longer time in the environment, also highlighting that in the cycle of this hemoparasite, transovarian transmission in *R. sanguineus* it is of great importance in the maintenance of this protozoan, causing ticks to remain infected for several generations²⁰. This information was corroborated by the results found in the present study, in which 41% (82/100) of the ticks were positive for *B. canis vogeli*.

No sample was positive for *R. vitalli*, even though a specimen of *Amblyomma aureolatum*, vector of the protozoan⁵ was collected in the present study, which suggests the need to monitor this pathogen in the city, as this infection can cause severe disease in domestic dogs²¹.

In the present study, positivity of 0.5% (1/200) of the samples for *E. canis* was detected through real-time PCR; 41% (82/100) for *B. canis vogeli*. The low percentage found for the bacterium stands out, as studies show the high prevalence of the disease in dogs in the country²². Probably due to the high infection found in these ticks for *B. canis vogeli*, could somehow inhibit the infection for *E. canis*, which would explain the difference in the results found; but further studies are needed to verify if the vector competence of the arthropod for these pathogens occurs in the same way and if co-infections in the vector may inhibit the proliferation of one of the pathogens.

It is important to emphasize that due to the nidicola behavior of *R. sanguineus* s.l., the environmental differences have little impact on the environmental infestation of this vector²³, sug-

gesting that sanitary management and responsible custody are important behaviors in the control of these hemoparasitoses.

CONCLUSION

Detection of *B. canis vogeli* and *E. canis* in ticks collected from dogs residing in the municipality of Itu, in the State of São Paulo, Brazil; show the dispersion of these pathogens in the country and; the role of *R. sanguineus* s.l. as a vector of these pathogens cannot be neglected.

ACKNOWLEDGMENT

Thanks to Fundação de Amparo à Pesquisa de São Paulo (FAPESP) for all financial support (case number 2016 / 00167-0).

REFERENCES

1. FONSECA, A. D. V., OLIVEIRA, L. M. B.; JORGE, F. R.; CAV-ALCANTE, R. O.; BEVILAQUA, C. M. L.; PINTO, F. J. M.; SANTOS, J. M. L.; TEIXEIRA, B. M.; RODRIGUES, A. K. P. P.; BRAZ, G. F.; VIANA, G. A.; COSTA, E. C.; SERPA, M. C. A.; WECK, B. C.; LABRUNA, M. B. Ocorrência de patógenos transmitidos por carrapatos em cães em uma região litorânea do estado do Ceará, nordeste do Brasil. *Revista Brasileira de Parasitologia Veterinária*, v. 31, n. 1, 2022.
2. OLIVER, J. H. Biology and systematics of ticks (Acari: Ixodidae). *Annual Review of Ecology and Systematics*, v. 20, p. 397-430, 1989.
3. MCDADE, J. E.; NEWHOUSE, V. F. Natural history of *Rickettsia rickettsii*. *Annu Ver Microbiol*, v. 40, p. 287-309, 1986.
4. MORAES-FILHO, J.; PINTER, A.; PACHECO, R. C.; GUTMANN, T. B.; BARBOSA, S. O.; GONZALES, M. A. R. M.; MURARO, M. A.; CECILIO, S. E. M.; LABRUNA, M. B. New Epidemiological Data on Brazilian Spotted Fever in an Endemic Area of the State of Sao Paulo, Brazil. *Vector Borne Zoonotic Dis*, v. 9, n. 1, p.73-78, 2009.
5. SOARES, J.F.; COSTA, F. B.; GIROTTO-SOARES, A.; DA SILVA, A.S.; FRANÇA, R.T.; TANIWAKI, S.A.; DALLAGNOL, B.; RECK, J.; HAGIWARA, M.K.; LABRUNA, M.B. Evaluation of the vector competence of six ixodid tick species for *Rangelia vitalli* (Apicomplexa, Piroplasmorida), the agent of canine rangeli-osis. *Ticks and Tick-Borne Diseases*, v. 1, p. 1-1234, 2018.
6. ZHANG, J.; LIU, Q.; WANG, D.; LI, W.; BEUGNET, F.; ZHOU, J. Epidemiological survey of ticks and tick-borne pathogens in pet dogs in south-eastern China. *Parasite*, v. 24, n. 35, p. 1-8, 2017.
7. DAGNONE, A. S.; MORAIS, H. S. A.; VIDOTTO, M. C.; JOJIMA, F. S.; VIDOTTO, O. Ehrlichiosis in anemic, thrombocytopenic, or tick-infested dogs from a hospital population in south Brazil. *Veterinary Parasitology*, v. 117, n. 4, p. 285-290, 2003.
8. JEFFERIES, R.; RYAN, U. M.; JARDINE, J.; BROUGHTON, D. K.; ROBERTSON, I. D.; IRWIN, P. J. Blood, bull terriers and babesiosis: further evidence for direct transmission of *Babesia gibsoni* in dogs. *Australian Veterinary Journal*, v.85, p.459-463, 2007.
9. IRWIN, Peter J. Canine babesiosis: from molecular taxonomy to control. *Parasites & vectors*, v. 2, n. 1, p. 1-9, 2009.
10. RAMOS, R.; RAMOS, C.; ARAUJO, F. R.; OLIVEIRA, R. H. M. de; SOUZA, I.; PIMENTEL, D.; GALINDO, M.; SANTAN, M.; ROSAS, E.; FAUSTINO, M.; ALVES, L. Molecular survey and genetic characterization of tick-borne pathogens in dogs in metropolitan Recife (north-eastern Brazil). *Parasitology research*, v. 107, n. 5, p. 1115-1120, 2010.
11. ZABÓ, M. P. J.; CUNHA, T. M.; PINTER, A.; VICENTINI, F. Ticks (Acari: Ixodidae) associated with domestic dogs in Franca region, São Paulo, Brazil. *Experimental and Applied Acarology*, v. 25, p. 909-916, 2001.
12. LABRUNA, M. B. & PEREIRA, M. C. Carrapato em cães no Brasil. *Clínica Veterinária*, v. 6, n. 30, p. 24-32, 2001.
13. SOARES, J. F., GIROTTO, A., BRANDÃO, P.E., SILVA, A.S., FRANÇA, R.T., LOPES, S.T.A., LABRUNA, M.B. 2011. Detection and molecular characterization of a canine piroplasm from Brazil. *Vet. Parasit.*, v.180, n.3, p.153-167, 2011.
14. BARROS-BATTESTI, D.M.; ARZUA, M.; BECHARA, G.H. Carrapatos de importância médico-veterinária da região neotropical: um guia ilustrado para identificação de espécies. São Paulo, Vox/ICTTD-3/Butantan. p. 223, 2006.
15. PAULINO, P.G.; PIRES, M.S.; DA SILVA C.B.; PECKLE, M., DA COSTA, R.L.; VITARI, G.L.V.; DE ABREU, A.P.M.; MASSARD, C.L.; SANTOS, H.A. Molecular epidemiology of *Babesia vogeli* in dogs from the southeastern region of Rio de Janeiro, Brazil. *Vet Parasitol Reg Stud Reports*, v. 13, p. 160-165, 2018.
16. DOYLE, C.K.; LABRUNA, M.B.; BREITSCHWERDT, E.B., TANG, Y.W., CORSTVET, R.E., HEGARTY, B.C., BLOCH, K.C., LI, P., WALKER, D.H., MCBRIDE, J.W. 2005. Detection of medically important Ehrlichia by quantitative multicolor TaqMan real-time polymerase chain reaction of the dsb gene. *J. Mol. Diagn.*, v.7, n.4, p.504-510, 2005.
17. SCINACHI, C. A.; TAKEDA, G. A. C. G.; MUCCI, L. F.; PINTER, A. Association of the occurrence of Brazilian spotted fever and Atlantic rain forest fragmentation in the São Paulo metropolitan region, Brazil. *Acta Tropica*, v. 166, p. 225-233, 2017.
18. ARAGÃO, H.; FONSECA, F. Notas de Ixodologia: IX. O complexo ovale do gênero *Amblyomma*. *Memórias do Instituto Oswaldo Cruz*, v. 59, n. 2, p. 131-148, 1961.
19. NEER, T.M. Canine monocytic and granulocytic ehrlichiosis. In: GREENE, C.E. Infectious diseases of dog and cat. Philadelphia: WB Saunders. 1998. p. 139-147.
20. MEHLHORN, H.; SHEIN, E. The piroplasms: life cycle and sexual stages. *Adv Parasitol.*, v.23, p. 37-103, 1984.
21. LEMOS, T. D.; TOMA, H. K.; ASSAD, R. Q.; SILVA, A. V.; CORRÊA, R. G. B.; ALMONSNY, N. R. P. 2017. Clinical and hematological evaluation of *Rangelia vitalli*-naturally infected dogs in southeastern Brazil. *Rev. Bras. Paras. Vet.*, v.26, p.307-313, 2017.
22. VIEIRA, R.F.C.; BIONDO, A.W.; GUIMARAES, A.M.S.; SANTOS, A.P.; SANTOS, R.P.; DUTRA, L.H.; DINIZ, P.P.V.P.; MORAIS, H.A.; MESSICK, J.B.; LABRUNA, M.B., VIDOTTO, O. Ehrlichiosis in Brazil. *Rev. Bras. Parasitol. Vet.*, v. 20, n. 1, p. 1-12, 2011.
23. DANTAS-TORRES, F. The brown dog tick, *Rhipicephalus sanguineus* (Latreille,1806) (Acari: Ixodidae): from taxonomy to control. *Vet Parasitol*, v.152:173-185, 2008.